

OTOSIGHT® CHARTING REMINDERS

Thorough and accurate charting is imperative for patients presenting with ear issues.

Diagnosis	<p>Use specific & accurate ICD-10-CM, for example:</p> <p>H65.03 – Acute serous otitis media, bilateral</p> <p>H66.006 – Acute suppurative otitis media without spontaneous rupture of the eardrum, recurrent, bilateral</p> <p>H73.013 – Bullous myringitis, bilateral</p> <p>H92.01 – Otalgia, right ear</p> <p>H92.12 – Otorrhea, left ear</p>
Imaging	Provide OtoSight image(s) report taken during patient visit, if available
Middle ear detail	<p>Share exactly what is seen, specifically mentioning use of OCT and/or OtoSight, for example:</p> <ul style="list-style-type: none"> • "Middle ear fluid/effusion present in both ears on OCT ear exam" • "No middle ear fluid/effusion present in unilateral right ear OCT exam"
Symptoms	<p>Be specific. The patient presents with:</p> <ul style="list-style-type: none"> • Ear pain • High temperature • Difficulty hearing • Discharge running out of the ear • A feeling of pressure or fullness inside the ear • Any other symptoms associated with ear infection <p>For those too young to communicate:</p> <ul style="list-style-type: none"> • Rub or pull at their ear(s) • Lack of reaction to some sounds • Irritable or restless • Any other symptoms associated with ear infection
Related History	Reference any previous antibiotic(s), surgery(ies), or any other intervention(s) the patient may have had related to the ear(s)
Medication	What medication(s) is the patient currently taking? What treatment have you prescribed for the patient today (if any)?
Other physicians	Include referring physician's patient chart (if available)
Plan of action	Describe what has been recommended to the patient for specific symptom(s)